

Analytic multiplicative cocycles over holomorphic dynamical systems

Tien-Cuong Dinh

May 17, 2008

Abstract

We prove some properties of analytic multiplicative and sub-multiplicative cocycles. The results allow to construct natural invariant analytic sets associated to complex dynamical systems.

AMS classification: 37F, 32H

Key-words: multiplicative cocycle, sub-multiplicative cocycle, exceptional set

1 Introduction

In this paper we study some sub-multiplicative cocycles over complex dynamical systems. The notion of sub-multiplicative cocycle is classical. It lies at the root of the definition of Lyapunov exponents. In [6] cocycles which are upper semi-continuous with respect to the analytic Zariski topology, were introduced and studied. Applications to holomorphic dynamics were subsequently given in [7, 8]. Our purpose here is to introduce a new approach closer in spirit to [3, 5]. The properties we obtain strengthen former results of Favre. In Section 3 below, we give an application to equidistribution problems of our main theorem, see also [3, 5].

Let X be an irreducible compact complex space of dimension k , not necessarily smooth. Let $f : X \rightarrow X$ be an open holomorphic map. Consider the dynamical system associated to f and define $f^n := f \circ \cdots \circ f$, n times, the iterate of order n of f .

Definition 1.1. A sequence of functions $\kappa_n : X \rightarrow [1, +\infty[$, $n \geq 0$, is an *analytic sub-multiplicative* (resp. *multiplicative*) *cocycle* if for any $n, m \geq 0$ and any $x \in X$, we have

1. κ_n is upper semi-continuous for the Zariski topology on X and $\min_X \kappa_n = 1$;
2. $\kappa_{n+m}(x) \leq \kappa_n(x)\kappa_m(f^n(x))$ (resp. $\kappa_{n+m}(x) = \kappa_n(x)\kappa_m(f^n(x))$).

The first property implies that κ_n is bounded from above and equal to 1 out of a finite or countable union of proper analytic subsets of X . Moreover, for any $\delta > 1$, $\{\kappa_n \geq \delta\}$ is a proper analytic subset of X and there is $\delta' < \delta$ such that $\{\kappa_n < \delta\}$ is contained in $\{\kappa_n < \delta'\}$. The second property defines classical multiplicative and sub-multiplicative cocycles, see [13]. It implies that if M is a constant such that $\kappa_1 \leq M$, then $\kappa_n \leq M^n$ for $n \geq 1$.

The new tool we introduce is to define for any analytic sub-multiplicative cocycle $(\kappa_n)_{n \geq 0}$, its extension to the negative integers:

$$\kappa_{-n}(x) := \max_{y \in f^{-n}(x)} \kappa_n(y).$$

Since $\{\kappa_{-n} \geq \delta\} = f^n\{\kappa_n \geq \delta\}$, the function κ_{-n} is upper semi-continuous in the Zariski sense. Here is our main result.

Theorem 1.2. *The sequence of functions $[\kappa_{-n}]^{1/n}$ converges pointwise to a function κ_- . Moreover, for every $\delta > 1$, the level set $\{\kappa_- \geq \delta\}$ is a proper analytic subset of X which is invariant under f and is contained in the orbit of $\{\kappa_n \geq \delta^n\}$ for every $n \geq 0$. In particular, κ_- is upper semi-continuous in the Zariski sense.*

Using the function κ_- , we obtain a more uniform version of [6, Th. 2.3.5] for sub-multiplicative cocycles, which is proved to be useful in applications, see Section 3 below.

Corollary 1.3. *If $\limsup[\kappa_n(x)]^{1/n} \geq \delta$, then x is sent by some f^n to a point in the invariant analytic set $\{\kappa_- \geq \delta\}$. If (κ_n) is multiplicative, then $[\kappa_n]^{1/n}$ converge pointwise to a function κ_+ such that $\kappa_+ \circ f = \kappa_+$.*

The new and useful property here is that the analytic set $\{\kappa_- \geq \delta\}$ does not depend on x . Note that one can extend the study to the case where $f : X \rightarrow X$ is a continuous open map on an irreducible Noetherian topological space of finite dimension, see [6, 11]. In particular, the above results hold for open real analytic maps and for open algebraic maps on quasi-projective varieties. We will discuss the case of dominant meromorphic maps on Kähler compact manifolds in Section 3.

Note also that the condition $\min \kappa_n = 1$ in Definition 1.1 can be replaced by $\kappa_n \geq c^n$ for some constant $c > 0$. Indeed, let Y be a compact analytic set and b a point in Y . Consider the map $f' : X \times Y \rightarrow X \times Y$ defined by $f'(x, y) := (f(x), y)$ and the cocycle $\kappa'_n(x, y) := 1$ if $y \neq b$ and $\kappa'_n(x, b) := 2^n c^{-n} \kappa_n(x)$. Theorem 1.2 and Corollary 1.3 can be applied to f' and κ'_n . We deduce analogous results for $\delta > \delta^* := \min \kappa_-$ (in Theorem 1.2 and Corollary 1.3 we have $\delta^* = 1$).

2 Asymptotic behavior of cocycles

In this section, we prove the results stated above. Observe that $(\kappa_{nl})_{n \geq 0}$ is an analytic (sub-)multiplicative cocycle for f^l with $l \geq 1$. We have the following elementary lemmas.

Lemma 2.1. *Let $l \geq 1$ be an integer. Then*

$$\limsup_{n \rightarrow \infty} [\kappa_{nl}]^{1/(nl)} = \limsup_{n \rightarrow \infty} [\kappa_n]^{1/n} \quad \text{and} \quad \liminf_{n \rightarrow \infty} [\kappa_{nl}]^{1/(nl)} = \liminf_{n \rightarrow \infty} [\kappa_n]^{1/n}.$$

Proof. If M is a constant such that $\kappa_s \leq M$ for any $0 \leq s \leq l$, then

$$\kappa_{m+s}(x) \leq \kappa_m(x) \kappa_s(f^m(x)) \leq M \kappa_m(x).$$

We deduce from these inequalities that

$$M^{-1} \kappa_{(n+1)l} \leq \kappa_{nl+s} \leq M \kappa_{nl}.$$

Hence

$$\limsup_{n \rightarrow \infty} [\kappa_{nl+s}]^{1/(nl+s)} = \limsup_{n \rightarrow \infty} [\kappa_{nl}]^{1/(nl)}$$

for every $0 \leq s \leq l-1$. The first identity in the lemma follows. The last one is obtained in the same way. $\square$

Lemma 2.2. *Let $l \geq 1$ be an integer. Then*

$$\limsup_{n \rightarrow \infty} [\kappa_{-nl}]^{1/(nl)} = \limsup_{n \rightarrow \infty} [\kappa_{-n}]^{1/n} \quad \text{and} \quad \liminf_{n \rightarrow \infty} [\kappa_{-nl}]^{1/(nl)} = \liminf_{n \rightarrow \infty} [\kappa_{-n}]^{1/n}.$$

Proof. As in Lemma 2.1, we get

$$\kappa_{m+s}(x) \leq \kappa_s(x) \kappa_m(f^s(x)) \leq M \kappa_m(f^s(x)). \quad (1)$$

Therefore, if $a \in X$ and $b \in f^{-m-s}(a)$ are such that $\kappa_{m+s}(b) = \kappa_{-m-s}(a)$, then

$$\kappa_{-m-s}(a) = \kappa_{m+s}(b) \leq M \kappa_m(f^s(b)) \leq M \max_{x \in f^{-m}(a)} \kappa_m(x) = M \kappa_{-m}(a).$$

We deduce that

$$M^{-1} \kappa_{-(n+1)l} \leq \kappa_{-nl-s} \leq M \kappa_{-nl}.$$

The lemma follows. $\square$

For any irreducible analytic subset Y of X , define

$$\kappa_n(Y) := \min_{x \in Y} \kappa_n(x) \quad \text{and} \quad \kappa_{-n}(Y) := \min_{x \in Y} \kappa_{-n}(x).$$

Since κ_n and κ_{-n} are upper semi-continuous for the Zariski topology, the previous minimums always exist. Moreover, we have $\kappa_n(Y) = \kappa_n(x)$ and $\kappa_{-n}(Y) = \kappa_{-n}(x)$ for $x \in Y$ outside a finite or countable union of proper analytic subsets of Y .

Lemma 2.3. *Let Y be a periodic irreducible analytic subset of X . Then $[\kappa_n(Y)]^{1/n}$ converge to a constant $\kappa_+(Y)$. Moreover, for every $m \geq 0$, we have $\kappa_+(f^m(Y)) = \kappa_+(Y)$ and $\liminf [\kappa_{-n}(Y)]^{1/n} \geq \kappa_+(Y)$. If $\kappa_+(Y) \geq \delta$ then for every $m \geq 0$, Y is contained in the orbit of $\{\kappa_m \geq \delta^m\}$.*

Proof. Assume that $f^l(Y) = Y$ with $l \geq 1$. Using properties of cocycles for a generic point in Y , we have

$$\kappa_{(m+n)l}(Y) \leq \kappa_{ml}(Y)\kappa_{nl}(f^{ml}(Y)) = \kappa_{ml}(Y)\kappa_{nl}(Y).$$

Hence, the sequence of $[\kappa_{nl}(Y)]^{1/(nl)}$ is decreasing and converges to some constant $\kappa_+(Y)$ when n tends to infinity. The first assertion follows from Lemma 2.1 applied to a generic point in Y . By definition of κ_{-n} , since $f^l(Y) = Y$, we have $\kappa_{-nl}(Y) \geq \kappa_{nl}(Y)$. This and Lemma 2.2 imply the assertion on κ_{-n} .

For the second assertion, applying (1) for a generic point in Y or $f(Y)$ yields

$$M^{-1}\kappa_{n+1}(Y) \leq \kappa_n(f(Y)) \leq M\kappa_{n-l+1}(f^l(Y)) = M\kappa_{n-l+1}(Y).$$

This and the first assertion imply that $[\kappa_n(f(Y))]^{1/n}$ converge to $\kappa_+(Y)$. It follows that $\kappa_+(f(Y)) = \kappa_+(Y)$ and then $\kappa_+(f^m(Y)) = \kappa_+(Y)$ for every $m \geq 0$.

Assume that $\kappa_+(Y) \geq \delta$ and that Y is not contained in the orbit of $\{\kappa_m \geq \delta^m\}$ for a fixed $m \geq 0$. Since Y is periodic, $f^i(Y)$ is not contained in $\{\kappa_m \geq \delta^m\}$ for every $i \geq 0$. We then deduce that $\kappa_m(f^i(Y)) \leq \delta_0^m$ for some constant $\delta_0 < \delta$. This implies

$$\kappa_{nm}(Y) \leq \prod_{i=0}^{n-1} \kappa_m(f^{im}(Y)) \leq \delta_0^{nm}$$

that contradicts the inequality $\kappa_+(Y) \geq \delta$. $\square$

We have the following proposition.

Proposition 2.4. *Let q be an integer with $0 \leq q \leq k-1$. Let Ω be a dense Zariski open subset of X such that $f^{-1}(\Omega) \subset \Omega$. Assume that $\{\kappa_{-n_0} \geq \delta^{n_0}\} \cap \Omega$ is of dimension at most equal to q for some $n_0 \geq 1$ and some $\delta > 1$. Then there is an integer $n_1 \geq 1$ and an invariant proper analytic subset Σ of X , possibly reducible, which is empty or of pure dimension q , such that*

1. Σ is contained in the orbit of $\{\kappa_n \geq \delta^n\}$ for every $n \geq 0$;
2. $\liminf[\kappa_n(x)]^{1/n} \geq \delta$ and $\liminf[\kappa_{-n}(x)]^{1/n} \geq \delta$ for $x \in \Sigma$;
3. $\{\kappa_{-n_1} \geq \delta^{n_1}\} \cap \Omega \setminus \Sigma$ is of dimension $\leq q-1$ if $q \geq 1$ and is empty if $q = 0$.

Proof. Let $\tilde{\Sigma}$ denote the union of the irreducible components of dimension q in the closure of $\{\kappa_{-n_0} \geq \delta^{n_0}\} \cap \Omega$. Recall that κ_{-n_0} is upper semi-continuous and that $\{\kappa_{-n_0} \geq \delta^{n_0}\} \cap \Omega$ is of dimension less than or equal to q . Hence, there is a real number $1 \leq \delta_0 < \delta$ satisfying $\kappa_{n_0}(V) \leq \delta_0^{n_0}$ for any irreducible analytic subset V of dimension $\geq q$ of X such that $f^{n_0}(V)$ intersects $\Omega \setminus \tilde{\Sigma}$. We will cover $\tilde{\Sigma}$ by three analytic sets Σ , Σ' and Σ'' .

Define Σ as the union of the orbits of periodic irreducible components Y of $\tilde{\Sigma}$ such that $\kappa_+(Y) \geq \delta$. So, Σ is invariant, i.e. $f(\Sigma) = \Sigma$. Note that Σ is not

necessarily contained in $\tilde{\Sigma}$. By Lemma 2.3, we also have $\kappa_+(f^i(Y)) \geq \delta$ for every $i \geq 0$ and Σ satisfies the first two properties in the proposition. Let Σ' be the union of the orbits of periodic irreducible components Y of $\tilde{\Sigma}$ with $\kappa_+(Y) < \delta$. Then, Σ' is invariant and by Lemma 2.3, $\kappa_+(Z) < \delta$ for any irreducible component Z of Σ' . Therefore, we can find $m \geq 0$ and $1 \leq \delta_1 < \delta$ such that $\kappa_{mn_0}(Z) \leq \delta_1^{mn_0}$ for any irreducible component Z of Σ' . Finally denote by Σ'' the union of the non-periodic irreducible components of $\tilde{\Sigma}$ and l the number of irreducible components of Σ'' .

Choose an integer N large enough so that $M^{l+m} \max(\delta_0, \delta_1)^{Nmn_0} < \delta^{Nmn_0}$ where M denotes the maximal value of κ_{n_0} on X . It is enough to check the last property in the proposition for $n_1 := Nmn_0$. For this purpose, we only have to show that $\kappa_{-n_1}(V_0) < \delta^{n_1}$ for any irreducible analytic set V_0 of dimension $q' \geq q$ which intersects $\Omega \setminus \Sigma$. Consider such an analytic set V_0 and an arbitrary sequence $V_{-Nm}, V_{-Nm+1}, \dots, V_0$ of irreducible analytic subsets of dimension q' of X such that $V_{-i+1} = f^{n_0}(V_{-i})$. Since $f^{-1}(\Omega \setminus \Sigma) \subset \Omega \setminus \Sigma$ and since f is surjective, all these analytic sets intersect $\Omega \setminus \Sigma$. It is also clear that the considered sequence contains at most l elements which are irreducible components of Σ'' . We show that $\kappa_{Nmn_0}(V_{-Nm}) < \delta^{Nmn_0}$ which implies that $\kappa_{-n_1}(V_0) < \delta^{n_1}$ and completes the proof. We distinguish three cases.

If V_0 is not an irreducible component of Σ' (in particular when $q' > q$), the sequence of V_{-n} does not contain any irreducible component of Σ' . It contains at most l elements which are irreducible components of Σ'' . The other components intersect $\Omega \setminus \tilde{\Sigma}$. Therefore, we have

$$\kappa_{Nmn_0}(V_{-Nm}) \leq \prod_{n=0}^{Nm-1} \kappa_{n_0}(V_{-Nm+n}) \leq M^l \delta_0^{Nmn_0} < \delta^{Nmn_0}.$$

From now on, we only have to consider the case where $q' = q$.

If V_{-Nm} is an irreducible component of Σ' then V_{-i} is an irreducible component of Σ' for $0 \leq i \leq Nm$ since Σ' is invariant. It follows from properties of cocycles that

$$\kappa_{Nmn_0}(V_{-Nm}) \leq \prod_{i=0}^{N-1} \kappa_{mn_0}(V_{-(N-i)m}) \leq \delta_1^{Nmn_0} < \delta^{Nmn_0}.$$

Otherwise, let $0 \leq s \leq N-1$ be the largest integer such that V_{-im} is an irreducible component of Σ' for $i \leq s$. Then V_{-n} is not an irreducible component of Σ' for $n \geq (s+1)m$. Therefore, since the considered sequence contains at most

l elements which are irreducible components of Σ'' , we have

$$\begin{aligned}
\kappa_{Nmn_0}(V_{-Nm}) &\leq \prod_{n=0}^{Nm-sm-1} \kappa_{n_0}(V_{-Nm+n}) \prod_{i=0}^{s-1} \kappa_{mn_0}(V_{-(s-i)m}) \\
&\leq \prod_{n=0}^{Nm-(s+1)m-1} \kappa_{n_0}(V_{-Nm+n}) \prod_{Nm-(s+1)m}^{Nm-sm-1} \kappa_{n_0}(V_{-Nm+n}) \prod_{i=0}^{s-1} \kappa_{mn_0}(V_{-(s-i)m}) \\
&\leq M^l \delta_0^{[Nm-(s+1)m]n_0} M^m \delta_1^{smn_0} < \delta^{Nmn_0}.
\end{aligned}$$

This completes the proof. $\square$

Proof of Theorem 1.2. Fix a real number $\delta > 1$. We apply Proposition 2.4 to $n_0 := 1$, $q := k - 1$ and $\Omega = \Omega_0 := X$. Since $\min \kappa_1 = 1$, we can find an invariant hypersurface Σ_1 and an integer n_1 such that $\{\kappa_{-n_1} \geq \delta^{n_1}\} \setminus \Sigma_1$ is of dimension at most equal to $k - 2$. Moreover, $\liminf[\kappa_n]^{1/n} \geq \delta$ and $\liminf[\kappa_{-n}]^{1/n} \geq \delta$ on Σ_1 . The Zariski open set $\Omega_1 := X \setminus \Sigma_1$ satisfies $f^{-1}(\Omega_1) \subset \Omega_1$. Hence, we can apply Proposition 2.4 inductively on q . We find invariant analytic subsets Σ_{k-q} of pure dimension q , $0 \leq q \leq k - 1$, such that $\liminf[\kappa_n]^{1/n} \geq \delta$ and $\liminf[\kappa_{-n}]^{1/n} \geq \delta$ on Σ_q . Moreover, if $\Omega_{k-q} := X \setminus \Sigma_1 \cup \dots \cup \Sigma_{k-q}$, then $\{\kappa_{-n_{k-q}} \geq \delta^{n_{k-q}}\} \cap \Omega_{k-q}$ is of dimension at most equal to $q - 1$ for some integer n_{k-q} .

When $q = 0$ the last intersection is empty. We deduce that $\kappa_{-n_k} < \delta_0^{n_k}$ on Ω_k for some constant $\delta_0 < \delta$. This, the fact that $f^{-1}(\Omega_k) \subset \Omega_k$ together with the properties of cocycles imply that $\kappa_{-mn_k} \leq \delta_0^{mn_k}$ on Ω_k for $m \geq 0$. By Lemma 2.2, $\limsup[\kappa_{-n}]^{1/n} \leq \delta_0$ on Ω_k .

If x is an arbitrary point in X , define $\kappa_{-}(x) := \liminf[\kappa_{-n}(x)]^{1/n}$. When $\kappa_{-}(x) < \delta$, x belongs to Ω_k . Hence, $\limsup[\kappa_{-n}(x)]^{1/n} \leq \delta_0 < \delta$. This holds for any $\delta > \kappa_{-}(x)$. We deduce that $[\kappa_{-n}(x)]^{1/n}$ converge to $\kappa_{-}(x)$. We also obtain from the previous analysis that $\{\kappa_{-} \geq \delta\}$ is equal to $\Sigma_1 \cup \dots \cup \Sigma_k$. By Proposition 2.4, this set is contained in the orbit of $\{\kappa_n \geq \delta^n\}$ for every $n \geq 0$. $\square$

Proof of Corollary 1.3. Let x be a point in X . Assume that $x_m := f^{mn_k}(x)$ is in Ω_k for every $m \geq 0$. We have $\kappa_{-n_k}(x_m) \leq \delta_0^{n_k}$ for every $m \geq 0$ and hence $\kappa_{n_k}(x_{m-1}) \leq \delta_0^{n_k}$ for every $m \geq 1$. Since (κ_n) is sub-multiplicative, $\kappa_{mn_k}(x) \leq \delta_0^{mn_k}$ for every $m \geq 1$. Lemma 2.1 implies that $\limsup[\kappa_n(x)]^{1/n} \leq \delta_0 < \delta$. The first assertion follows.

Now assume that (κ_n) is multiplicative. Define $\kappa_{+}(x) := \liminf[\kappa_n(x)]^{1/n}$. We have seen that if x belongs to $\Sigma_1 \cup \dots \cup \Sigma_k$ then $\kappa_{+}(x) \geq \delta$. So, the properties of multiplicative cocycles imply that if x is sent by a f^n to a point in $\Sigma_1 \cup \dots \cup \Sigma_k$ then $\kappa_{+}(x) \geq \delta$. In other words, for any $\delta > \kappa_{+}(x)$ and $m \geq 0$, $x_m := f^{mn_k}(x)$ is in Ω_k . Therefore, we have $\limsup[\kappa_n(x)]^{1/n} \leq \delta_0 < \delta$. This implies that $\limsup[\kappa_n(x)]^{1/n} \leq \kappa_{+}(x)$. Hence, $[\kappa_n(x)]^{1/n}$ converge to $\kappa_{+}(x)$. The identity $\kappa_{+} \circ f = \kappa_{+}$ is a consequence of the properties of multiplicative cocycles. $\square$

Proposition 2.5. *Let (κ_n) be a sub-multiplicative cocycle as above. Let x be a point in X . Then, for $l \geq 0$ large enough, $[\kappa_n(f^l(x))]^{1/n}$ converge to a constant $\kappa(x)$ which does not depend on l . In particular, x is sent by an iterate of f to the invariant analytic set $\{\kappa_- \geq \kappa(x)\}$.*

Proof. Let $\chi_l := \limsup_{n \rightarrow \infty} [\kappa_n(f^l(x))]^{1/n}$. We have $\chi_l \geq 1$. Inequalities (1) for $s = 1$ imply that $\kappa_{n+1}(f^l(x)) \leq M\kappa_n(f^{l+1}(x))$. Therefore, the sequence $(\chi_l)_{l \geq 0}$ is increasing. Define $\delta := \lim_{l \rightarrow \infty} \chi_l = \sup_l \chi_l$. If $\delta = 1$, we have $\chi_l = 1$; hence $[\kappa_n(f^l(x))]^{1/n}$ converge to 1 for every $l \geq 0$. Assume now that $\delta > 1$. We use the notations and the arguments in the proofs of Theorem 1.2 and Corollary 1.3. When l is large enough, we have $\chi_l > \delta_0$. Therefore, $f^l(x)$ is sent by an iterate of f to a point in $\Sigma_1 \cup \dots \cup \Sigma_k$. So, for l large enough, $f^l(x)$ belongs to $\Sigma_1 \cup \dots \cup \Sigma_k$. For such an l , we have $\liminf_{n \rightarrow \infty} [\kappa_n(f^l(x))]^{1/n} \geq \delta$. This implies that $[\kappa_n(f^l(x))]^{1/n}$ converge to δ . The last assertion in the proposition is deduced from Corollary 1.3. $\square$

The following example shows that $[\kappa_n]^{1/n}$ do not always converge when n goes to infinity.

Example 2.6. Let a be a fixed point of f and b a point in $f^{-1}(a) \setminus \{a\}$. Let λ_n be positive numbers such that $\lambda_{n+1} \leq \lambda_n + 1$. Define the analytic sub-multiplicative cocycle (κ_n) by $\kappa_n(a) := e^n$, $\kappa_n(b) := e^{\lambda_n}$ and $\kappa_n(x) := 1$ for $x \neq a, b$. Then, $[\kappa_n(b)]^{1/n}$ converge if and only if the arguments of the complex numbers $n + i\lambda_n$ converge. It is not difficult to find λ_n which do not satisfy this condition. In this case, the cocycle (κ_n) is not multiplicative. When $\lambda_n = 0$, $[\kappa_n]^{1/n}$ converge but the limit is not an invariant function.

3 Exceptional sets for meromorphic maps

There are some natural cocycles over a complex dynamical system. We give some examples. Let $f : \mathbb{P}^k \rightarrow \mathbb{P}^k$ be a holomorphic map of algebraic degree $d \geq 2$ on the projective space of dimension k . It is induced by a map $F : \mathbb{C}^{k+1} \rightarrow \mathbb{C}^{k+1}$ whose irreducible components are homogeneous polynomials of degree d . The map f defines a ramified covering of degree d^k on $\mathbb{P}^k$. We refer to the survey [12] for the basic properties of f .

Let $\kappa_n(x)$ denote the multiplicity of f^n at x , i.e. $f^{-n}(z)$ contains exactly $\kappa_n(x)$ points near x for any z generic close enough to $f^n(x)$. It is not difficult to check that (κ_n) is an analytic multiplicative cocycle. We now give an application of Theorem 1.2.

Theorem 3.1 ([3, 5]). *Let $f : \mathbb{P}^k \rightarrow \mathbb{P}^k$ be a holomorphic map of algebraic degree $d \geq 2$. Then there is a proper analytic subset $\mathcal{E}$ of $\mathbb{P}^k$, possibly empty, which is totally invariant by f : $f^{-1}(\mathcal{E}) = f(\mathcal{E}) = \mathcal{E}$, and is maximal: $\mathcal{E}$ contains all the proper analytic subsets E of $\mathbb{P}^k$ satisfying $f^{-m}(E) \subset E$ for some $m \geq 1$.*

Proof. We use the multiplicative cocycle (κ_n) defined above. Let E be as in the theorem. The sequence of analytic sets $(f^{-nm}(E))$ is decreasing. Hence, there is an n such that $f^{-nm-m}(E) = f^{-nm}(E)$. Since f^{nm} is surjective, we deduce that $f^{-m}(E) = E$ and $f^m(E) = E$. Observe also that if $\tilde{E} := E \cup \dots \cup f^{-m+1}(E)$ then $f^{-1}(\tilde{E}) = \tilde{E}$. Replacing E by $\tilde{E}$ allows to assume that $m = 1$ and that E is totally invariant: $f^{-1}(E) = f(E) = E$. The union of the irreducible components of E , of a given fixed dimension, is also totally invariant. So, it is enough to consider the case where E is of pure dimension p . Let $n_0 \geq 1$ be an integer such that f^{n_0} fixes all the irreducible components of E .

Denote by g the restriction of f^{n_0} to E' , an irreducible component of E . We claim that the topological degree of g is equal to d^{pn_0} , that is, g defines a ramified covering of degree d^{pn_0} . Indeed, if ω is a Kähler form on $\mathbb{P}^k$, then $(f^{n_0})^*(\omega^p)$ is cohomologous to $d^{pn_0}\omega^p$. By Bézout theorem, the mass of the measure $g^*(\omega_{|E'}^p) = (f^{n_0})^*(\omega^p)|_{E'}$ is equal to d^{pn_0} times the mass of $\omega_{|E'}^p$. This implies the claim.

Now, observe that the topological degree of f^{n_0} is d^{kn_0} while the topological degree of g is equal to d^{pn_0} . This implies that $\kappa_{-n_0} \geq d^{(k-p)n_0}$ on E' since $f^{-n_0}(E') = E'$. The inequality still holds if we replace n_0 by a multiple of n_0 . Therefore, E' is contained in $\{\kappa_- \geq d^{k-p}\}$ and E is contained in $\mathcal{E}' := \{\kappa_- \geq d\}$. Since E is totally invariant, it is contained in $\mathcal{E} := \bigcap_{n \geq 0} f^{-n}(\mathcal{E}')$. As above, we prove that $\mathcal{E}$ is totally invariant. This completes the proof. $\square$

The above result still holds for the restriction of f to an invariant irreducible analytic set X of dimension p . One lifts f to a normalization of X . The map we obtain is a ramified covering of degree d^p . The cocycle (κ_n) over this map can be defined as above. A simple decreasing induction on dimension allows to prove that f admits a finite number of totally invariant analytic subsets. Moreover, the convex set of totally invariant probability measures, i.e. probability measures ν such that $f^*(\nu) = d^k\nu$, is of finite dimension. The *equilibrium measure* μ is the only totally invariant probability measure which has no mass on $\mathcal{E}$. The exceptional set $\mathcal{E}$ is also characterized by the property that $x \notin \mathcal{E}$ if and only if $d^{-kn}(f^n)^*(\delta_x)$ converge to μ [3, 5], see also [1, 9]. Here, δ_x denotes the Dirac mass at x .

Consider now a more general case. Let (X, ω) be a compact Kähler manifold of dimension k and let $f : X \rightarrow X$ be a dominant meromorphic map. So, f is a holomorphic map from $X \setminus I$ to X where I is the set of indeterminacy which is analytic and of codimension ≥ 2 . The closure Γ of the graph of $f : X \setminus I \rightarrow X$ in $X \times X$ is an analytic set. If $\pi_i : X \times X \rightarrow X$, $i = 1, 2$, are the natural projections, then f is equal to $\pi_2 \circ (\pi_1|_{\Gamma})^{-1}$ on $X \setminus I$. Define for $A \subset X$

$$f(A) := \pi_2(\pi_1^{-1}(A) \cap \Gamma) \quad \text{and} \quad f^{-1}(A) := \pi_1(\pi_2^{-1}(A) \cap \Gamma).$$

Define also $f^n := f \circ \dots \circ f$, n times, on a suitable dense Zariski open set and extend it to a meromorphic map on X . In general, we do not have $f^2(A) = f(f(A))$ nor $f^{-2}(A) = f^{-1}(f^{-1}(A))$.

The set of indeterminacy I is the set of points x such that $f(x)$ is of dimension at least equal to 1. Define $I_1 := I$ and $I_{n+1} := f^{-1}(I_n)$ for $n \geq 1$. The orbit of a point x

$$x, f(x), f^2(x), \dots, f^n(x), \dots$$

is well-defined if $x \notin \cup_{n \geq 1} I_n$.

Let I' denote the set of points x such that $f^{-1}(x)$ is of dimension larger than or equal to 1. This is also an analytic subset of codimension ≥ 2 of X . The fibers $f^{-1}(x)$ for $x \notin I'$, are finite and contain the same number of points counted with multiplicity. This number d_t is called *topological degree* of f . If ν is a finite measure on X having no mass on I' , then $f^*(\nu)$ is well-defined by the formula

$$f^*(\nu) := (\pi_1)_*((\pi_2|_\Gamma)^*\nu).$$

If ν is positive, the mass of $f^*(\nu)$ is equal to d_t times the mass of ν . Define $I'_1 := I'$ and $I'_{n+1} := f(I'_n)$ for $n \geq 1$. If $x \notin \cup_{n \geq 1} I'_n$, then $f^{-n}(x)$ contains exactly d_t^n points counting with multiplicity.

Choose a finite or countable union $\tilde{I}$ of proper analytic subsets of X containing $I \cup I'$ such that if $\tilde{X} := X \setminus \tilde{I}$ then $f^{-1}(\tilde{X}) = f(\tilde{X}) = \tilde{X}$. So, $f : \tilde{X} \rightarrow \tilde{X}$ defines a “ramified covering” of degree d_t . We consider the topology on $\tilde{X}$ induced by the Zariski topology on X . As in the case of holomorphic maps on X , one can define cocycles over $f : \tilde{X} \rightarrow \tilde{X}$. Theorem 1.2 and Corollary 1.3 still hold in this case.

Recall that the *dynamical degree* of order p of f , $0 \leq p \leq k$, is defined by

$$d_p := \lim_{n \rightarrow \infty} \left(\int_X (f^n)^*(\omega^p) \wedge \omega^{k-p} \right)^{1/n}.$$

Note that $(f^n)^*(\omega^p)$ is a (p, p) -form with L^1 coefficients. The last limit always exists and the dynamical degrees are bi-meromorphic invariants [4]. We also have $d_0 = 1$ and $d_k = d_t$, the topological degree of f .

Theorem 3.2. *Let $f : X \rightarrow X$ be a dominant meromorphic map as above. Assume that the topological degree of f is strictly larger than the other dynamical degrees. Then there is a proper analytic subset $\mathcal{E}$ of X , possibly empty, such that*

1. *all the irreducible components of $\mathcal{E}$ intersect $\tilde{X}$;*
2. *$\mathcal{E}$ is totally invariant: $f^{-1}(\mathcal{E} \cap \tilde{X}) = f(\mathcal{E} \cap \tilde{X}) = \mathcal{E} \cap \tilde{X}$;*
3. *$\mathcal{E}$ is maximal: if $E \subset X$ is a proper analytic subset whose irreducible components intersect $\tilde{X}$ and if $f^{-m}(E \cap \tilde{X}) \subset E$ for some $m \geq 1$, then $E \subset \mathcal{E}$.*

One can also characterize $\mathcal{E}$ by the following property. For x a point in $\tilde{X}$, the sequence of probability measures $d_t^{-n}(f^n)^*(\delta_x)$ converges to the equilibrium measure of f if and only if $x \notin \mathcal{E}$, see [2, 5, 10] for the proof.

The proof of Theorem 3.2 follows the arguments in Theorem 3.1. We only need the following lemma.

Lemma 3.3. *Let E be a proper irreducible analytic subset of X which intersects $\tilde{X}$ and $n_0 \geq 1$ an integer such that $f^{n_0}(E \cap \tilde{X}) \subset E$. Then the topological degree of the meromorphic map $f^{n_0} : E \rightarrow E$ is at most equal to δ^{n_0} where $0 < \delta < d_t$ is a constant independent of E and n_0 .*

Proof. Denote by g the map $f^{n_0} : E \rightarrow E$. Let p be the dimension of E , d the topological degree of g and $\delta < d_t$ a constant larger than or equal to the dynamical degrees $d_0, \dots, d_{k-1}$. The mass of $g^*(\omega|_E^p)$ is equal to d times the mass of $\omega|_E^p$. We will bound the mass of $g^*(\omega|_E^p)$. Fix a constant $\epsilon > 0$. In what follows, all the constants are independent of E , n_0 and n .

The form $(f^{n_0})^*(\omega^p)$ has L^1 coefficients and is smooth on the Zariski open set $X \setminus \cup_{1 \leq n \leq n_0} I_n$ which contains $\tilde{X}$. It also defines a positive closed (p, p) -current on X whose mass is given by the formula

$$\|(f^{n_0})^*(\omega^p)\| := \int_X (f^{n_0})^*(\omega^p) \wedge \omega^{k-p}.$$

Therefore, there is a constant $c > 0$ such that this mass is bounded by $c(d_p + \epsilon)^{n_0}$. We regularize $(f^{n_0})^*(\omega^p)$ using the results in [4]. There are smooth positive closed (p, p) -forms $\Omega_i^\pm$ such that $\Omega_i^+ - \Omega_i^-$ converge to $(f^{n_0})^*(\omega^p)$ in the sense of currents and

$$\|\Omega_i^\pm\| \leq A\|(f^{n_0})^*(\omega^p)\| \leq A'(d_p + \epsilon)^{n_0}$$

for some constants A and A' . Since $(f^{n_0})^*(\omega^p)$ is smooth in a neighbourhood of $\tilde{X}$, the construction in [4] gives forms $\Omega_i^\pm$ such that $\Omega_i^+ - \Omega_i^-$ converge uniformly on compact subsets of $\tilde{X}$ to $(f^{n_0})^*(\omega^p)$. It follows that

$$\|g^*(\omega|_E^p)\| = \|(f^{n_0})^*(\omega^p)|_{E \cap \tilde{X}}\| \leq \limsup_{i \rightarrow \infty} \int_E \Omega_i^+.$$

The last integral depends only on the cohomology class of Ω_i^+ . The above bound on the mass of Ω_i^+ induces a bound on its cohomology class. Therefore, $\|g^*(\omega|_E^p)\| \leq A''(d_p + \epsilon)^{n_0}$ for some constant A'' . The estimate also holds for the iterates of g : we have $\|(g^n)^*(\omega|_E^p)\| \leq A''(d_p + \epsilon)^{nn_0}$. It follows that $d^n \lesssim (d_p + \epsilon)^{nn_0}$. Hence, d is at most equal to $d_p^{n_0}$. This completes the proof. $\square$

References

- [1] Briend J.-Y., Duval J., Deux caractérisations de la mesure d'équilibre d'un endomorphisme de $P^k(\mathbf{C})$, *Publ. Math. Inst. Hautes Études Sci.*, **93** (2001), 145-159.
- [2] Dinh T.-C., Suites d'applications méromorphes multivaluées et courants laminaires, *J. Geom. Anal.*, **15** (2005), no. 2, 207-227.

- [3] Dinh T.-C., Sibony N., Dynamique des applications d'allure polynomiale, *J. Math. Pures Appl.*, **82** (2003), 367-423.
- [4] ———, Regularization of currents and entropy, *Ann. Sci. École Norm. Sup.*, **37** (2004), 959-971.
- [5] ———, Equidistribution towards the Green current for holomorphic maps, *Ann. Sci. École Norm. Sup.*, to appear.
- [6] Favre C., Dynamique des applications rationnelles, *PhD thesis*, Université Paris-Sud, Orsay, 2000.
- [7] ———, Multiplicity of holomorphic functions, *Math. Ann.*, **316** (2000), no. 2, 355-378.
- [8] Favre C., Jonsson M., Brolin's theorem for curves in two complex dimensions, *Ann. Inst. Fourier*, **53** (2003), no. 5, 1461-1501.
- [9] Fornæss J.-E., Sibony N., Complex dynamics in higher dimension. II. Modern methods in complex analysis (Princeton, NJ, 1992), 135-182, *Ann. of Math. Stud.*, **137**, Princeton Univ. Press, Princeton, NJ, 1995.
- [10] Guedj V., Ergodic properties of rational mappings with large topological degree, *Ann. of Math. (2)*, **161** (2005), no. 3, 1589-1607.
- [11] Hartshorne R., *Algebraic geometry*, Graduate Texts in Mathematics, No. **52**, Springer-Verlag, New York-Heidelberg, 1977.
- [12] Sibony N., Dynamique des applications rationnelles de $\mathbb{P}^k$, *Panoramas et Synthèses*, **8** (1999), 97-185.
- [13] Walters P., *An introduction to ergodic theory*, Graduate Texts in Mathematics, **79**, Springer-Verlag, New York-Berlin, 1982.

T.-C. Dinh, UPMC Univ Paris 06, UMR 7586, Institut de Mathématiques de Jussieu, F-75005 Paris, France. dinh@math.jussieu.fr, <http://www.math.jussieu.fr/~dinh>